

Pan-Ox-5™

A complete dietary enzyme formula

DESCRIPTION

Pan-Ox-5™, provided by Douglas Laboratories, is a synergistic dietary enzyme formula that is specially formulated for digestive support.

FUNCTIONS

The pancreas supplies the major digestive enzymes that catalyze the breakdown of starches (carbohydrates), proteins, and fats, so that the breakdown products can be absorbed in the upper small intestine.

Some carbohydrate digestion takes place in the mouth by salivary amylase, but pancreatic amylase is the major carbohydrate-digesting enzyme. Amylases break down starches to maltose and maltotriose, which are further hydrolyzed into glucose by the disaccharidases of the mucosal cells, and then absorbed. Although cellulose is indigestible by pancreatic and intestinal enzymes, the microflora within the large intestine may degrade it and produce uncomfortable symptoms in the process.

Although fat digestion starts in the mouth with the action of salivary lipase, the great majority of fat triglycerides are digested by pancreatic lipase secreted by the exocrine pancreas into the duodenum. Lipases break down triglycerides into monoglycerides and free fatty acids, which are efficiently absorbed in the upper small intestine.

Protein digestion is initiated in the stomach by pepsin and hydrochloric acid, which denature and break large proteins down to smaller polypeptides. In the small intestine, proteases break down these polypeptides into free amino acids, and di- and tripeptides, which are directly absorbed by the intestinal mucosa.

Ingredients such as aspergillus oryza, ox bile extract and rennin can also aid in the digestion and absorption of food.

INDICATIONS

Pan-Ox-5™ may be a useful nutritional adjunct for individuals who wish to supplement their diets with a wide variety of digestive enzymes.

FORMULA (#7917)

Each Tablet Contains:

Betaine HCl	60 mg
Glutamic Acid	10 mg
Pancreatin (4X).....	75 mg
(equal to 300 mg of Pancreatin)	
Pepsin (1:3000).....	32.5 mg
Papain	48 mg
Aspergillus Oryza	35 mg
Lipase	50 mg
Ox Bile Extract	65 mg
Rennin	15 mg
Malt Diastase	7.5 mg
Beet Root Powder	100 mg
Citrus Pectin	12.5 mg

In a base of Myrrh Gum.

SUGGESTED USE

Adults take 1 tablet daily with meals or as directed by physician.

SIDE EFFECTS

No adverse side effects reported.

STORAGE

Store in a cool, dry place, away from direct light. Keep out of reach of children.

REFERENCES

Braden B, Picard H, Caspary WF, et al. Monitoring pancreatic supplementation in cystic fibrosis patients with the ¹³C-Hiolein breath test: evidence for normalized fat assimilation with high dose pancreatic therapy. *Z Gastroenterol* 1997;35:123-9.

(continued on reverse)

Bragelmann R, Armbrecht U, Rosemeyer D, et al. The effect of pancreatic enzyme supplementation in patients with steatorrhea after total gastrectomy [see comments]. *Eur J Gastroenterol Hepatol* 1999;11:231-7.

Gan KH, Heijerman HG, Geus WP, et al. Comparison of a high lipase pancreatic enzyme extract with a regular pancreatin preparation in adult cystic fibrosis patients. *Aliment Pharmacol Ther* 1994;8:603-7.

Greenberger NJ. Enzymatic therapy in patients with chronic pancreatitis. *Gastroenterol Clin North Am* 1999;28:687-93.

Kalivianakis M, Minich DM, Bijleveld CM, et al. Fat malabsorption in cystic fibrosis patients receiving enzyme replacement therapy is due to impaired intestinal uptake of long-chain fatty acids. *Am J Clin Nutr* 1999;69:127-34.

Layer P, Keller J. Pancreatic enzymes: secretion and luminal nutrient digestion in health and disease. *J Clin Gastroenterol*

1999;28:3-10.

Neoptolemos JP, Ghaneh P, Andren-Sandberg A, et al. Treatment of pancreatic exocrine insufficiency after pancreatic resection. Results of a randomized, double-blind, placebo-controlled, crossover study of high vs standard dose pancreatin. *Int J Pancreatol* 1999;25:171-80.

Suarez F, Levitt MD, Adshead J, et al. Pancreatic supplements reduce symptomatic response of healthy subjects to a high fat meal. *Dig Dis Sci* 1999;44:1317-21.

Taylor CJ, Hillel PG, Ghosal S, et al. Gastric emptying and intestinal transit of pancreatic enzyme supplements in cystic fibrosis. *Arch Dis Child* 1999;80:149-52.

Van Hoozen CM, Peeke PG, Taubeneck M, et al. Efficacy of enzyme supplementation after surgery for chronic pancreatitis. *Pancreas* 1997;14:174-80.

**These statements have not been evaluated by the Food and Drug Administration.
This product is not intended to diagnose, treat, cure, or prevent any disease.**

**Manufactured by
Douglas Laboratories
600 Boyce Road
Pittsburgh, PA 15205
800-245-4440**